

LAKEWATCH Report for Blackwater in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Blackwater River		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Blackwater
Latitude	25.9343
Longitude	-81.5956
Water Body Type	Estuary
Period of Record (year)	2001 to 2001

LAKEWATCH Report for Blackwater in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	63 - 63	63 (1)
Total Nitrogen (µg/L)	494 - 494	494 (1)
Chlorophyll- uncorrected (µg/L)	5.7 - 5.7	5.7 (1)
Secchi (ft)	2.3 - 2.3	2.3 (1)
Secchi (m)	0.7 - 0.7	0.7 (1)
Color (Pt-Co Units)	18 -18	18 (1)
Specific Conductance (µS/cm@25 C)	51500 - 51500	51500 (1)
Salinity (ppt)	32 - 32	32 (1)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll (µg/L) (Forsberg and Ryding 1980)	Coastal Chlorophyll (µg/L) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 – 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Blackwater in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This values ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Blackwater in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Cape Romano in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Rookery Bay Marco Island		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Cape Romano
Latitude	25.8569
Longitude	-81.6726
Water Body Type	Estuary
Period of Record (year)	2001 to 2004

LAKEWATCH Report for Cape Romano in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	23 - 31	27 (4)
Total Nitrogen ($\mu\text{g/L}$)	113 - 318	232 (4)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	1.3 - 4.0	2.9 (4)
Secchi (ft)	1.8 - 7.0	4.2 (4)
Secchi (m)	0.5 - 2.1	1.3 (4)
Color (Pt-Co Units)	3 -4	4 (2)
Specific Conductance ($\mu\text{S/cm@25 C}$)	49000 - 54000	51500 (2)
Salinity (ppt)	31 - 34	32 (2)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Cape Romano in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This values ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Cape Romano in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Clam Bay East-1 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Clam Bay (Collier County)	Clam Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Clam Bay East-1
Latitude	26.2311
Longitude	-81.8129
Water Body Type	Estuary
Period of Record (year)	2001 to 2003

LAKEWATCH Report for Clam Bay East-1 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	389 - 449	411 (3)
Total Nitrogen ($\mu\text{g/L}$)	2689 - 3355	3073 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	35.8 - 39.0	37.2 (3)
Secchi (ft)	0.7 - 1.0	0.8 (3)
Secchi (m)	0.2 - 0.3	0.2 (3)
Color (Pt-Co Units)	148 - 189	167 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	1000 - 1500	1278 (3)
Salinity (ppt)	0 - 1	0 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Clam Bay East-1 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This values ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Clam Bay East-1 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Clam Bay East-2 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Clam Bay (Collier County)	Clam Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Clam Bay East-2
Latitude	26.2416
Longitude	-81.8161
Water Body Type	Estuary
Period of Record (year)	2001 to 2003

LAKEWATCH Report for Clam Bay East-2 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	99 - 121	109 (3)
Total Nitrogen (µg/L)	903 - 999	960 (3)
Chlorophyll- uncorrected (µg/L)	16.3 - 18.0	17.4 (3)
Secchi (ft)	1.4 - 2.0	1.6 (3)
Secchi (m)	0.4 - 0.6	0.5 (3)
Color (Pt-Co Units)	49 -84	65 (3)
Specific Conductance (µS/cm@25 C)	25500 - 41250	33583 (3)
Salinity (ppt)	16 - 26	21 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll (µg/L) (Forsberg and Ryding 1980)	Coastal Chlorophyll (µg/L) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 – 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Clam Bay East-2 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This values ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Clam Bay East-2 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Clam Bay East-3 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Clam Bay (Collier County)	Clam Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Clam Bay East-3
Latitude	26.2385
Longitude	-81.8171
Water Body Type	Estuary
Period of Record (year)	2001 to 2003

LAKEWATCH Report for Clam Bay East-3 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	109 - 125	116 (3)
Total Nitrogen ($\mu\text{g/L}$)	886 - 1018	968 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	14.8 - 18.3	15.9 (3)
Secchi (ft)	1.4 - 2.2	1.7 (3)
Secchi (m)	0.4 - 0.7	0.5 (3)
Color (Pt-Co Units)	49 -69	59 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	27000 - 42750	33472 (3)
Salinity (ppt)	17 - 27	21 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Clam Bay East-3 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This values ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Clam Bay East-3 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Clam Bay West-1 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Clam Bay (Collier County)	Clam Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Clam Bay West-1
Latitude	26.2439
Longitude	-81.8206
Water Body Type	Estuary
Period of Record (year)	2001 to 2003

LAKEWATCH Report for Clam Bay West-1 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	588 - 1376	905 (3)
Total Nitrogen ($\mu\text{g/L}$)	3292 - 6850	4839 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	22.4 - 65.2	49.3 (3)
Secchi (ft)	0.1 - 0.7	0.5 (3)
Secchi (m)	0.0 - 0.2	0.1 (3)
Color (Pt-Co Units)	231 -383	305 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	12000 - 12500	12250 (3)
Salinity (ppt)	7 - 8	7 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Clam Bay West-1 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This values ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Clam Bay West-1 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Clam Bay West-2 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Clam Bay (Collier County)	Clam Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Clam Bay West-2
Latitude	26.2375
Longitude	-81.8129
Water Body Type	Estuary
Period of Record (year)	2001 to 2003

LAKEWATCH Report for Clam Bay West-2 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	131 - 324	206 (3)
Total Nitrogen ($\mu\text{g/L}$)	1192 - 2195	1727 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	16.6 - 28.1	23.9 (3)
Secchi (ft)	0.3 - 1.7	1.0 (3)
Secchi (m)	0.1 - 0.5	0.3 (3)
Color (Pt-Co Units)	94 -141	125 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	34250 - 48250	39278 (3)
Salinity (ppt)	21 - 30	24 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Clam Bay West-2 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This values ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Clam Bay West-2 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Clam Bay West-3 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Clam Bay (Collier County)	Clam Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Clam Bay West-3
Latitude	26.223
Longitude	-81.8167
Water Body Type	Estuary
Period of Record (year)	2001 to 2003

LAKEWATCH Report for Clam Bay West-3 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	57 - 69	63 (3)
Total Nitrogen ($\mu\text{g/L}$)	505 - 529	516 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	3.5 - 9.3	6.7 (3)
Secchi (ft)	1.2 - 1.7	1.5 (3)
Secchi (m)	0.4 - 0.5	0.5 (3)
Color (Pt-Co Units)	8 -25	18 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	36250 - 52500	43806 (3)
Salinity (ppt)	23 - 33	27 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Clam Bay West-3 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This values ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Clam Bay West-3 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Fakahatchee Bay in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Gulf Islands		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Fakahatchee Bay
Latitude	25.8922
Longitude	-81.477
Water Body Type	Estuary
Period of Record (year)	2001 to 2001

LAKEWATCH Report for Fakahatchee Bay in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	54 - 54	54 (1)
Total Nitrogen ($\mu\text{g/L}$)	524 - 524	524 (1)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	6.4 - 6.4	6.4 (1)
Secchi (ft)	2.5 - 2.5	2.5 (1)
Secchi (m)	0.8 - 0.8	0.8 (1)
Color (Pt-Co Units)	18 - 18	18 (1)
Specific Conductance ($\mu\text{S/cm@25 C}$)	47500 - 47500	47500 (1)
Salinity (ppt)	30 - 30	30 (1)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Fakahatchee Bay in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Fakahatchee Bay in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Fakaunion in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Gulf Islands		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Fakaunion
Latitude	25.9014
Longitude	-81.5159
Water Body Type	Estuary
Period of Record (year)	2001 to 2001

LAKEWATCH Report for Fakaunion in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	32 - 32	32 (1)
Total Nitrogen ($\mu\text{g/L}$)	371 - 371	371 (1)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	4.3 - 4.3	4.3 (1)
Secchi (ft)	2.9 - 2.9	2.9 (1)
Secchi (m)	0.9 - 0.9	0.9 (1)
Color (Pt-Co Units)	11 -11	11 (1)
Specific Conductance ($\mu\text{S/cm@25 C}$)	45500 - 45500	45500 (1)
Salinity (ppt)	28 - 28	28 (1)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Fakaunion in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Fakaunion in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Johnson Bay 1 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Rookery Bay Marco Island		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Johnson Bay 1
Latitude	25.9858
Longitude	-81.7256
Water Body Type	Estuary
Period of Record (year)	2001 to 2011

LAKEWATCH Report for Johnson Bay 1 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	25 - 41	34 (11)
Total Nitrogen ($\mu\text{g/L}$)	195 - 366	301 (11)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	3.0 - 9.0	4.9 (11)
Secchi (ft)	3.0 - 4.8	4.0 (11)
Secchi (m)	0.9 - 1.5	1.2 (11)
Color (Pt-Co Units)	5 - 9	7 (10)
Specific Conductance ($\mu\text{S/cm@25 C}$)	46667 - 52200	49939 (10)
Salinity (ppt)	29 - 33	31 (10)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Johnson Bay 1 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	11	11	11	11
Intercept (a)	735	-15600	205	-58
Slope (b)	-0.35	7.93	-0.10	0.03
Coefficient of Determination (R ²)	0.06	0.28	0.04	0.03
Probability of Significance (p)	0.47	0.10	0.55	0.61
Potential Trend	No Trend	No Trend	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Johnson Bay 1 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Johnson Bay 1 (Collier)




Johnson Bay 1 (Collier)



Johnson Bay 1 (Collier)



Johnson Bay 1 (Collier)



LAKEWATCH Report for Johnson Bay 2 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Rookery Bay Marco Island		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Johnson Bay 2
Latitude	25.9918
Longitude	-81.7207
Water Body Type	Estuary
Period of Record (year)	2001 to 2011

LAKEWATCH Report for Johnson Bay 2 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	30 - 43	36 (11)
Total Nitrogen ($\mu\text{g/L}$)	186 - 376	312 (11)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	3.2 - 8.5	5.1 (11)
Secchi (ft)	1.8 - 5.0	3.3 (11)
Secchi (m)	0.5 - 1.5	1.0 (11)
Color (Pt-Co Units)	5 -16	9 (10)
Specific Conductance ($\mu\text{S/cm@25 C}$)	46333 - 52000	48936 (10)
Salinity (ppt)	29 - 32	30 (10)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Johnson Bay 2 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	11	11	11	11
Intercept (a)	814	-14459	309	95
Slope (b)	-0.39	7.36	-0.15	-0.05
Coefficient of Determination (R ²)	0.11	0.17	0.08	0.02
Probability of Significance (p)	0.32	0.21	0.40	0.68
Potential Trend	No Trend	No Trend	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Johnson Bay 2 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Johnson Bay 2 (Collier)




Johnson Bay 2 (Collier)



Johnson Bay 2 (Collier)



Johnson Bay 2 (Collier)



LAKEWATCH Report for Johnson Bay 3 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Rookery Bay Marco Island		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Johnson Bay 3
Latitude	25.9796
Longitude	-81.732
Water Body Type	Estuary
Period of Record (year)	2004 to 2011

LAKEWATCH Report for Johnson Bay 3 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	25 - 39	32 (8)
Total Nitrogen ($\mu\text{g/L}$)	209 - 368	306 (8)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	3.2 - 7.8	5.3 (8)
Secchi (ft)	3.8 - 5.5	4.6 (8)
Secchi (m)	1.1 - 1.7	1.4 (8)
Color (Pt-Co Units)	5 - 9	7 (7)
Specific Conductance ($\mu\text{S/cm@25 C}$)	41833 - 52667	49236 (7)
Salinity (ppt)	26 - 33	31 (7)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Johnson Bay 3 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	8	8	8	8
Intercept (a)	548	-29580	-341	466
Slope (b)	-0.26	14.89	0.17	-0.23
Coefficient of Determination (R ²)	0.03	0.46	0.07	0.67
Probability of Significance (p)	0.70	0.07	0.52	0.01
Potential Trend	No Trend	No Trend	No Trend	Decreasing

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Johnson Bay 3 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Johnson Bay 3 (Collier)




Johnson Bay 3 (Collier)



Johnson Bay 3 (Collier)



Johnson Bay 3 (Collier)



LAKEWATCH Report for Naples Bay in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Naples Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay
Latitude	26.1298
Longitude	-81.7921
Water Body Type	Estuary
Period of Record (year)	2001 to 2001

LAKEWATCH Report for Naples Bay in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	61 - 61	61 (1)
Total Nitrogen ($\mu\text{g/L}$)	507 - 507	507 (1)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	8.8 - 8.8	8.8 (1)
Secchi (ft)	3.1 - 3.1	3.1 (1)
Secchi (m)	0.9 - 0.9	0.9 (1)
Color (Pt-Co Units)	13 -13	13 (1)
Specific Conductance ($\mu\text{S/cm@25 C}$)	46000 - 46000	46000 (1)
Salinity (ppt)	29 - 29	29 (1)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Naples Bay-AQS 8 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Naples Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-AQS 8
Latitude	26.1293
Longitude	-81.8012
Water Body Type	Estuary
Period of Record (year)	2004 to 2006

LAKEWATCH Report for Naples Bay-AQS 8 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	62 - 100	74 (3)
Total Nitrogen ($\mu\text{g/L}$)	490 - 693	597 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	11.0 - 14.8	12.7 (3)
Secchi (ft)	3.4 - 3.8	3.6 (3)
Secchi (m)	1.0 - 1.2	1.1 (3)
Color (Pt-Co Units)	14 -38	28 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	23000 - 36000	31000 (3)
Salinity (ppt)	14 - 22	19 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-AQS 8 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-AQS 8 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Naples Bay-ARS 896 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-ARS 896
Latitude	26.2114
Longitude	-81.7682
Water Body Type	Estuary
Period of Record (year)	2004 to 2016

LAKEWATCH Report for Naples Bay-ARS 896 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	22 - 63	39 (13)
Total Nitrogen ($\mu\text{g/L}$)	590 - 931	766 (13)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	2.6 - 18.2	11.0 (13)
Secchi (ft)	2.8 - 4.4	3.4 (13)
Secchi (m)	0.9 - 1.4	1.0 (13)
Color (Pt-Co Units)	11 -33	22 (13)
Specific Conductance ($\mu\text{S/cm@25 C}$)	384 - 1000	572 (13)
Salinity (ppt)	0 - 0	0 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-ARS 896 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	13	13	13	13
Intercept (a)	2457	2993	-703	140
Slope (b)	-1.20	-1.11	0.36	-0.07
Coefficient of Determination (R ²)	0.16	0.00	0.06	0.29
Probability of Significance (p)	0.17	0.90	0.42	0.06
Potential Trend	No Trend	No Trend	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-ARS 896 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Naples Bay-ARS 896 (Collier)




Naples Bay-ARS 896 (Collier)



Naples Bay-ARS 896 (Collier)



Naples Bay-ARS 896 (Collier)



LAKEWATCH Report for Naples Bay-DOLLAR 15 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Rookery Bay Marco Island		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-DOLLAR 15
Latitude	26.0895
Longitude	-81.7868
Water Body Type	Estuary
Period of Record (year)	2004 to 2006

LAKEWATCH Report for Naples Bay-DOLLAR 15 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	38 - 79	53 (3)
Total Nitrogen ($\mu\text{g/L}$)	327 - 435	394 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	5.4 - 7.0	6.3 (3)
Secchi (ft)	2.6 - 4.7	3.8 (3)
Secchi (m)	0.8 - 1.4	1.2 (3)
Color (Pt-Co Units)	5 -12	8 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	47000 - 50500	48722 (3)
Salinity (ppt)	29 - 31	30 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-DOLLAR 15 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-DOLLAR 15 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Naples Bay-GORD 10 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida’s waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida’s waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria’s for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Naples Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-GORD 10
Latitude	26.093
Longitude	-81.7986
Water Body Type	Estuary
Period of Record (year)	2004 to 2006

LAKEWATCH Report for Naples Bay-GORD 10 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	38 - 59	46 (3)
Total Nitrogen ($\mu\text{g/L}$)	276 - 355	314 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	4.3 - 6.3	5.5 (3)
Secchi (ft)	2.5 - 6.0	4.3 (3)
Secchi (m)	0.8 - 1.8	1.3 (3)
Color (Pt-Co Units)	6 -9	8 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	49000 - 50000	49667 (3)
Salinity (ppt)	31 - 31	31 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-GORD 10 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-GORD 10 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Naples Bay-GORD 70 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Naples Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-GORD 70
Latitude	26.152
Longitude	-81.7856
Water Body Type	Estuary
Period of Record (year)	2004 to 2006

LAKEWATCH Report for Naples Bay-GORD 70 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	59 - 84	71 (3)
Total Nitrogen ($\mu\text{g/L}$)	598 - 715	647 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	8.7 - 17.0	11.9 (3)
Secchi (ft)	2.8 - 4.3	3.6 (3)
Secchi (m)	0.8 - 1.3	1.1 (3)
Color (Pt-Co Units)	22 -45	31 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	19300 - 29000	25100 (3)
Salinity (ppt)	12 - 18	15 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-GORD 70 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-GORD 70 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Naples Bay-GORD 80 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Naples Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-GORD 80
Latitude	26.1599
Longitude	-81.7838
Water Body Type	Estuary
Period of Record (year)	2004 to 2006

LAKEWATCH Report for Naples Bay-GORD 80 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	47 - 61	56 (3)
Total Nitrogen ($\mu\text{g/L}$)	560 - 698	641 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	6.5 - 18.0	11.0 (3)
Secchi (ft)	3.0 - 3.6	3.3 (3)
Secchi (m)	0.9 - 1.1	1.0 (3)
Color (Pt-Co Units)	30 -48	37 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	15000 - 20000	18322 (3)
Salinity (ppt)	9 - 12	11 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-GORD 80 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-GORD 80 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Naples Bay-GRE 896 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-GRE 896
Latitude	26.1738
Longitude	-81.7846
Water Body Type	Estuary
Period of Record (year)	2004 to 2016

LAKEWATCH Report for Naples Bay-GRE 896 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	25 - 62	37 (13)
Total Nitrogen ($\mu\text{g/L}$)	536 - 814	686 (13)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	6.4 - 20.6	11.7 (13)
Secchi (ft)	0.9 - 2.4	1.9 (13)
Secchi (m)	0.3 - 0.7	0.6 (13)
Color (Pt-Co Units)	18 -29	23 (13)
Specific Conductance ($\mu\text{S/cm@25 C}$)	346 - 1000	553 (13)
Salinity (ppt)	0 - 0	0 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-GRE 896 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	13	13	13	13
Intercept (a)	2981	13486	516	-10
Slope (b)	-1.46	-6.37	-0.25	0.01
Coefficient of Determination (R ²)	0.32	0.08	0.06	0.00
Probability of Significance (p)	0.04	0.35	0.42	0.85
Potential Trend	Decreasing	No Trend	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-GRE 896 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Naples Bay-GRE 896 (Collier)




Naples Bay-GRE 896 (Collier)



Naples Bay-GRE 896 (Collier)



Naples Bay-GRE 896 (Collier)



LAKEWATCH Report for Naples Bay-NBAY 22 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Naples Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-NBAY 22
Latitude	26.1095
Longitude	-81.7922
Water Body Type	Estuary
Period of Record (year)	2004 to 2006

LAKEWATCH Report for Naples Bay-NBAY 22 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	44 - 50	47 (3)
Total Nitrogen ($\mu\text{g/L}$)	350 - 462	412 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	7.3 - 10.5	8.4 (3)
Secchi (ft)	2.9 - 4.3	3.6 (3)
Secchi (m)	0.9 - 1.3	1.1 (3)
Color (Pt-Co Units)	8 -12	11 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	45667 - 50000	47889 (3)
Salinity (ppt)	28 - 31	30 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-NBAY 22 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-NBAY 22 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Naples Bay-NBAY 24 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Naples Bay		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Naples Bay-NBAY 24
Latitude	26.1132
Longitude	-81.786
Water Body Type	Estuary
Period of Record (year)	2004 to 2006

LAKEWATCH Report for Naples Bay-NBAY 24 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	42 - 72	53 (3)
Total Nitrogen ($\mu\text{g/L}$)	335 - 446	392 (3)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	5.7 - 9.3	7.1 (3)
Secchi (ft)	2.0 - 3.9	3.1 (3)
Secchi (m)	0.6 - 1.2	0.9 (3)
Color (Pt-Co Units)	6 -14	9 (3)
Specific Conductance ($\mu\text{S/cm@25 C}$)	44333 - 49500	47278 (3)
Salinity (ppt)	28 - 31	29 (3)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Naples Bay-NBAY 24 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Naples Bay-NBAY 24 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Pumpkin Bay in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Blackwater River		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Pumpkin Bay
Latitude	25.9176
Longitude	-81.5484
Water Body Type	Estuary
Period of Record (year)	2001 to 2001

LAKEWATCH Report for Pumpkin Bay in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	52 - 52	52 (1)
Total Nitrogen ($\mu\text{g/L}$)	439 - 439	439 (1)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	11.6 - 11.6	11.6 (1)
Secchi (ft)	2.3 - 2.3	2.3 (1)
Secchi (m)	0.7 - 0.7	0.7 (1)
Color (Pt-Co Units)	10 -10	10 (1)
Specific Conductance ($\mu\text{S/cm@25 C}$)	52000 - 52000	52000 (1)
Salinity (ppt)	32 - 32	32 (1)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Pumpkin Bay in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Pumpkin Bay in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Tarpon Bay 1 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Rookery Bay Marco Island		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Tarpon Bay 1
Latitude	25.986
Longitude	-81.7241
Water Body Type	Estuary
Period of Record (year)	2001 to 2011

LAKEWATCH Report for Tarpon Bay 1 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	35 - 53	43 (11)
Total Nitrogen ($\mu\text{g/L}$)	333 - 500	415 (11)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	3.0 - 13.8	9.2 (11)
Secchi (ft)	3.1 - 7.5	4.6 (11)
Secchi (m)	0.9 - 2.3	1.4 (11)
Color (Pt-Co Units)	9 -13	10 (9)
Specific Conductance ($\mu\text{S/cm@25 C}$)	43000 - 53833	50369 (9)
Salinity (ppt)	27 - 34	31 (9)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Tarpon Bay 1 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	11	11	11	11
Intercept (a)	-676	-13888	-926	396
Slope (b)	0.36	7.13	0.47	-0.19
Coefficient of Determination (R ²)	0.04	0.18	0.21	0.29
Probability of Significance (p)	0.55	0.20	0.15	0.11
Potential Trend	No Trend	No Trend	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Tarpon Bay 1 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Tarpon Bay 1 (Collier)




Tarpon Bay 1 (Collier)



Tarpon Bay 1 (Collier)



Tarpon Bay 1 (Collier)



LAKEWATCH Report for Tarpon Bay 2 in Collier County Using Data Downloaded 10/17/2016

Introduction Estuary

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Estuaries and coastal segments: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).

The near shore Florida coastline is separated into estuary and estuary segments within the estuary. Deeper coastal waters are separated into coastal nutrient regions and coastal nutrient segments within the regions. Numeric nutrient criteria are established for all estuary segments, including criteria for total nitrogen, total phosphorus, and chlorophyll a. For open ocean coastal waters, numeric criteria are established for chlorophyll a, that is derived from satellite remote sensing techniques. For those locations without defined segments there are narrative nutrient criteria (e.g., Florida Keys Halo Zone).

The maps defining individual estuaries and coastal segments can be found at: <https://www.flrules.org/Gateway/reference.asp?No=Ref-05420>.

The individual nutrient criteria can be found at: <https://www.flrules.org/gateway/ruleNo.asp?id=62-302.532>

Estuary lies in the following location:

Estuary	Estuary Segment	Coastal Nutrient Region	Coastal Nutrient Segment
Tidal Cocohatchee River/Ten Thousands Islands	Rookery Bay Marco Island		

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the estuary resides.
- **Name:** Estuary name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years an estuary has been in the LAKEWATCH program.

County	Collier
Name	Tarpon Bay 2
Latitude	25.9786
Longitude	-81.7295
Water Body Type	Estuary
Period of Record (year)	2004 to 2011

LAKEWATCH Report for Tarpon Bay 2 in Collier County Using Data Downloaded 10/17/2016

Long-Term Data Summary Estuary: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus ($\mu\text{g/L}$):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen ($\mu\text{g/L}$):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected ($\mu\text{g/L}$):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance ($\mu\text{S/cm@25}^\circ\text{C}$), Salinity (ppt):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Estuary: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus ($\mu\text{g/L}$)	27 - 40	34 (8)
Total Nitrogen ($\mu\text{g/L}$)	250 - 403	332 (8)
Chlorophyll- uncorrected ($\mu\text{g/L}$)	4.3 - 7.0	5.5 (8)
Secchi (ft)	3.3 - 5.9	4.5 (8)
Secchi (m)	1.0 - 1.8	1.4 (8)
Color (Pt-Co Units)	6 -9	7 (7)
Specific Conductance ($\mu\text{S/cm@25 C}$)	43500 - 55167	49815 (7)
Salinity (ppt)	27 - 34	31 (7)

Coastal Trophic State

Trophic status is a measure of a systems biological productivity and LAKEWATCH uses total chlorophyll averages as a trophic state measure. Since the total chlorophyll measurement indicates how much algae is actually present in a water body, it is the most direct indicator of biological productivity. For freshwater lakes, LAKEWATCH uses the trophic state classification criteria proposed by Forsberg and Ryding (1980). LAKEWATCH staff sampled coastal systems around all of Florida (Hoyer et al. 2002) and discovered that chlorophyll concentrations are significantly less for the same amount of algae than freshwater lakes. Thus, to classify trophic status of coastal waters using similar classification terminology LAKEWATCH provided the table below accounting for the chlorophyll differences reported by Hoyer et al. (2002).

Trophic Status	Freshwater Chlorophyll ($\mu\text{g/L}$) (Forsberg and Ryding 1980)	Coastal Chlorophyll ($\mu\text{g/L}$) (Hoyer et al. 2002)
Oligotrophic	< 3.0	< 0.5
Mesotrophic	3.0 - 7.0	0.5 - 1.8
Eutrophic	7.0 - 40.0	1.8 - 12.4
Hypereutrophic	> 40.0	> 12.4

Hoyer, M. V., T. K. Frazer, S. K. Notestein and D. E. Canfield, Jr. 2002. Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. Canadian Journal of Fisheries and Aquatic Sciences 59:1-8.

LAKEWATCH Report for Tarpon Bay 2 in Collier County Using Data Downloaded 10/17/2016

Trend Analyses Estuary


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would cross if the x-axis were zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	8	8	8	8
Intercept (a)	-1150	-24025	-176	638
Slope (b)	0.59	12.13	0.09	-0.32
Coefficient of Determination (R ²)	0.08	0.40	0.06	0.71
Probability of Significance (p)	0.50	0.09	0.57	0.01
Potential Trend	No Trend	No Trend	No Trend	Decreasing

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Tarpon Bay 2 in Collier County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Tarpon Bay 2 (Collier)



Tarpon Bay 2 (Collier)



Tarpon Bay 2 (Collier)



Tarpon Bay 2 (Collier)

