

LAKEWATCH Report for Black Creek-1 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	Black Creek-1
Latitude	30.451
Longitude	-86.07
Water Body Type	Stream
Period of Record (year)	1999 to 2000
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for Black Creek-1 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	12 - 13	12 (2)
Total Nitrogen (µg/L)	294 - 350	322 (2)
Chlorophyll- uncorrected (µg/L)	2.0 - 4.6	3.3 (2)
Secchi (ft)	4.5 - 5.0	4.8 (2)
Secchi (m)	1.4 - 1.5	1.4 (2)
Color (Pt-Co Units)	-	()
Specific Conductance (µS/cm@25 C)	-	()

LAKEWATCH Report for Black Creek-1 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Black Creek-1 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Black Creek-2 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	Black Creek-2
Latitude	30.4478
Longitude	-86.0795
Water Body Type	Stream
Period of Record (year)	1999 to 2000
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for Black Creek-2 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	9 - 12	11 (2)
Total Nitrogen (µg/L)	300 - 340	320 (2)
Chlorophyll- uncorrected (µg/L)	1.0 - 4.8	2.9 (2)
Secchi (ft)	4.3 - 5.0	4.6 (2)
Secchi (m)	1.3 - 1.5	1.4 (2)
Color (Pt-Co Units)	-	()
Specific Conductance (µS/cm@25 C)	-	()

LAKEWATCH Report for Black Creek-2 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Black Creek-2 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Black Creek-3 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	Black Creek-3
Latitude	30.4481
Longitude	-86.0919
Water Body Type	Stream
Period of Record (year)	1999 to 2000
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for Black Creek-3 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	12 - 13	13 (2)
Total Nitrogen (µg/L)	322 - 340	331 (2)
Chlorophyll- uncorrected (µg/L)	2.0 - 5.0	3.5 (2)
Secchi (ft)	3.9 - 3.9	3.9 (2)
Secchi (m)	1.2 - 1.2	1.2 (2)
Color (Pt-Co Units)	-	()
Specific Conductance (µS/cm@25 C)	-	()

LAKEWATCH Report for Black Creek-3 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Black Creek-3 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA Baker Canal-1 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA Baker Canal-1
Latitude	30.3229
Longitude	-86.1436
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA Baker Canal-1 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida’s fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	29 - 29	29 (1)
Total Nitrogen (µg/L)	500 - 500	500 (1)
Chlorophyll- uncorrected (µg/L)	13.5 - 13.5	13.5 (1)
Secchi (ft)	2.7 - 2.7	2.7 (1)
Secchi (m)	0.8 - 0.8	0.8 (1)
Color (Pt-Co Units)	19 - 19	19 (1)
Specific Conductance (µS/cm@25 C)	11500 - 11500	11500 (1)

LAKEWATCH Report for CBA Baker Canal-1 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA Baker Canal-1 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA Baker Canal-2 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).



Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA Baker Canal-2
Latitude	30.3755
Longitude	-86.155
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA Baker Canal-2 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	30 - 30	30 (1)
Total Nitrogen (µg/L)	453 - 453	453 (1)
Chlorophyll- uncorrected (µg/L)	13.0 - 13.0	13.0 (1)
Secchi (ft)	2.4 - 2.4	2.4 (1)
Secchi (m)	0.7 - 0.7	0.7 (1)
Color (Pt-Co Units)	17 - 17	17 (1)
Specific Conductance (µS/cm@25 C)	9500 - 9500	9500 (1)

LAKEWATCH Report for CBA Baker Canal-2 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA Baker Canal-2 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA Little Canal-1 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA Little Canal-1
Latitude	30.3737
Longitude	-86.1588
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA Little Canal-1 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	24 - 24	24 (1)
Total Nitrogen (µg/L)	438 - 438	438 (1)
Chlorophyll- uncorrected (µg/L)	10.8 - 10.8	10.8 (1)
Secchi (ft)	2.9 - 2.9	2.9 (1)
Secchi (m)	0.9 - 0.9	0.9 (1)
Color (Pt-Co Units)	34 - 34	34 (1)
Specific Conductance (µS/cm@25 C)	6835 - 6835	6835 (1)

LAKEWATCH Report for CBA Little Canal-1 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA Little Canal-1 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA Little Canal-2 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA Little Canal-2
Latitude	30.378
Longitude	-86.1613
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA Little Canal-2 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	25 - 25	25 (1)
Total Nitrogen (µg/L)	423 - 423	423 (1)
Chlorophyll- uncorrected (µg/L)	9.5 - 9.5	9.5 (1)
Secchi (ft)	2.9 - 2.9	2.9 (1)
Secchi (m)	0.9 - 0.9	0.9 (1)
Color (Pt-Co Units)	22 - 22	22 (1)
Specific Conductance (µS/cm@25 C)	9580 - 9580	9580 (1)

LAKEWATCH Report for CBA Little Canal-2 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA Little Canal-2 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-1 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-1
Latitude	30.4299
Longitude	-86.1134
Water Body Type	Stream
Period of Record (year)	2003 to 2005
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-1 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	25 - 47	33 (3)
Total Nitrogen (µg/L)	423 - 546	478 (3)
Chlorophyll- uncorrected (µg/L)	3.6 - 5.8	4.9 (3)
Secchi (ft)	2.4 - 3.0	2.6 (3)
Secchi (m)	0.7 - 0.9	0.8 (3)
Color (Pt-Co Units)	39 - 58	52 (3)
Specific Conductance (µS/cm@25 C)	900 - 1000	947 (3)

LAKEWATCH Report for CBA River-1 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-1 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-10 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-10
Latitude	30.5593
Longitude	-85.9578
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-10 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	6 - 6	6 (1)
Total Nitrogen (µg/L)	497 - 497	497 (1)
Chlorophyll- uncorrected (µg/L)	0.7 - 0.7	0.7 (1)
Secchi (ft)	-	()
Secchi (m)	-	()
Color (Pt-Co Units)	69 - 69	69 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-10 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-10 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-11 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).



Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-11
Latitude	30.4518
Longitude	-85.9356
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-11 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	12 - 12	12 (1)
Total Nitrogen (µg/L)	1100 - 1100	1100 (1)
Chlorophyll- uncorrected (µg/L)	1.0 - 1.0	1.0 (1)
Secchi (ft)	2.0 - 2.0	2.0 (1)
Secchi (m)	0.6 - 0.6	0.6 (1)
Color (Pt-Co Units)	296 - 296	296 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-11 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-11 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-12 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-12
Latitude	30.4664
Longitude	-85.9006
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-12 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	12 - 12	12 (1)
Total Nitrogen (µg/L)	1240 - 1240	1240 (1)
Chlorophyll- uncorrected (µg/L)	1.0 - 1.0	1.0 (1)
Secchi (ft)	-	()
Secchi (m)	-	()
Color (Pt-Co Units)	195 - 195	195 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-12 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-12 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-13 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).



Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-13
Latitude	30.4695
Longitude	-85.9981
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-13 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	8 - 8	8 (1)
Total Nitrogen (µg/L)	700 - 700	700 (1)
Chlorophyll- uncorrected (µg/L)	1.0 - 1.0	1.0 (1)
Secchi (ft)	-	()
Secchi (m)	-	()
Color (Pt-Co Units)	158 - 158	158 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-13 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-13 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-14 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).



Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-14
Latitude	30.5576
Longitude	-85.9604
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-14 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	6 - 6	6 (1)
Total Nitrogen (µg/L)	520 - 520	520 (1)
Chlorophyll- uncorrected (µg/L)	1.0 - 1.0	1.0 (1)
Secchi (ft)	-	()
Secchi (m)	-	()
Color (Pt-Co Units)	69 - 69	69 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-14 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-14 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-2 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-2
Latitude	30.4413
Longitude	-85.9618
Water Body Type	Stream
Period of Record (year)	2003 to 2004
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-2 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	26 - 29	28 (2)
Total Nitrogen (µg/L)	527 - 552	539 (2)
Chlorophyll- uncorrected (µg/L)	1.3 - 2.8	2.1 (2)
Secchi (ft)	1.8 - 2.7	2.2 (2)
Secchi (m)	0.5 - 0.8	0.7 (2)
Color (Pt-Co Units)	10 - 40	25 (2)
Specific Conductance (µS/cm@25 C)	900 - 900	900 (2)

LAKEWATCH Report for CBA River-2 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-2 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-3 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-3
Latitude	30.4512
Longitude	-85.898
Water Body Type	Stream
Period of Record (year)	2010 to 2016
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-3 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	11 - 30	26 (7)
Total Nitrogen (µg/L)	350 - 713	580 (7)
Chlorophyll- uncorrected (µg/L)	1.7 - 3.3	2.3 (7)
Secchi (ft)	2.0 - 3.5	2.8 (7)
Secchi (m)	0.6 - 1.1	0.8 (7)
Color (Pt-Co Units)	10 - 34	20 (7)
Specific Conductance (µS/cm@25 C)	57 - 137	103 (7)

LAKEWATCH Report for CBA River-3 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	7	7	7	7
Intercept (a)	-3051	-79804	105	38
Slope (b)	1.53	39.93	-0.05	-0.02
Coefficient of Determination (R ²)	0.25	0.60	0.03	0.00
Probability of Significance (p)	0.26	0.04	0.70	0.88
Potential Trend	No Trend	Increasing	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-3 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


CBA River-3 (Walton)




CBA River-3 (Walton)



CBA River-3 (Walton)



CBA River-3 (Walton)



LAKEWATCH Report for CBA River-3A in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-3A
Latitude	30.457
Longitude	-85.8977
Water Body Type	Stream
Period of Record (year)	2004 to 2010
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-3A in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	26 - 38	30 (7)
Total Nitrogen (µg/L)	480 - 842	632 (7)
Chlorophyll- uncorrected (µg/L)	2.0 - 5.1	3.3 (7)
Secchi (ft)	2.3 - 3.1	2.6 (7)
Secchi (m)	0.7 - 0.9	0.8 (7)
Color (Pt-Co Units)	20 - 37	25 (6)
Specific Conductance (µS/cm@25 C)	64 - 900	364 (6)

LAKEWATCH Report for CBA River-3A in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	7	7	7	7
Intercept (a)	-240	-94428	-983	64
Slope (b)	0.13	47.36	0.49	-0.03
Coefficient of Determination (R ²)	0.01	0.55	0.89	0.05
Probability of Significance (p)	0.88	0.06	0.00	0.64
Potential Trend	No Trend	No Trend	Increasing	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-3A in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


CBA River-3A (Walton)




CBA River-3A (Walton)



CBA River-3A (Walton)



CBA River-3A (Walton)



LAKEWATCH Report for CBA River-4 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).



Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-4
Latitude	30.4514
Longitude	-86.0689
Water Body Type	Stream
Period of Record (year)	2006 to 2016
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-4 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	13 - 19	16 (11)
Total Nitrogen (µg/L)	316 - 538	462 (11)
Chlorophyll- uncorrected (µg/L)	3.0 - 19.0	7.2 (11)
Secchi (ft)	3.0 - 4.7	3.7 (11)
Secchi (m)	0.9 - 1.4	1.1 (11)
Color (Pt-Co Units)	60 - 128	88 (10)
Specific Conductance (µS/cm@25 C)	41 - 4538	952 (10)

LAKEWATCH Report for CBA River-4 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	11	11	11	11
Intercept (a)	125	-20421	1450	63
Slope (b)	-0.05	10.38	-0.72	-0.03
Coefficient of Determination (R ²)	0.01	0.24	0.25	0.05
Probability of Significance (p)	0.80	0.13	0.12	0.50
Potential Trend	No Trend	No Trend	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-4 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


CBA River-4 (Walton)




CBA River-4 (Walton)



CBA River-4 (Walton)



CBA River-4 (Walton)



LAKEWATCH Report for CBA River-5 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).



Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-5
Latitude	30.5393
Longitude	-85.9205
Water Body Type	Stream
Period of Record (year)	2007 to 2016
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-5 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	5 - 11	7 (10)
Total Nitrogen (µg/L)	290 - 490	377 (10)
Chlorophyll- uncorrected (µg/L)	0.5 - 1.1	0.7 (10)
Secchi (ft)	2.5 - 4.5	3.1 (10)
Secchi (m)	0.8 - 1.4	1.0 (10)
Color (Pt-Co Units)	21 - 54	33 (10)
Specific Conductance (µS/cm@25 C)	18 - 392	77 (10)

LAKEWATCH Report for CBA River-5 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	10	10	10	10
Intercept (a)	-828	27353	14	939
Slope (b)	0.42	-13.41	-0.01	-0.47
Coefficient of Determination (R ²)	0.40	0.43	0.01	0.80
Probability of Significance (p)	0.05	0.04	0.81	0.04
Potential Trend	No Trend	Decreasing	No Trend	Decreasing

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-5 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


CBA River-5 (Walton)




CBA River-5 (Walton)



CBA River-5 (Walton)



CBA River-5 (Walton)



LAKEWATCH Report for CBA River-6 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-6
Latitude	30.4727
Longitude	-85.8884
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-6 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	25 - 25	25 (1)
Total Nitrogen (µg/L)	592 - 592	592 (1)
Chlorophyll- uncorrected (µg/L)	7.0 - 7.0	7.0 (1)
Secchi (ft)	2.8 - 2.8	2.8 (1)
Secchi (m)	0.9 - 0.9	0.9 (1)
Color (Pt-Co Units)	15 - 15	15 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-6 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-6 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-7 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-7
Latitude	30.45
Longitude	-85.9359
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-7 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	16 - 16	16 (1)
Total Nitrogen (µg/L)	808 - 808	808 (1)
Chlorophyll- uncorrected (µg/L)	2.5 - 2.5	2.5 (1)
Secchi (ft)	-	()
Secchi (m)	-	()
Color (Pt-Co Units)	289 - 289	289 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-7 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-7 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-8 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).



Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-8
Latitude	30.4655
Longitude	-85.8994
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-8 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	19 - 19	19 (1)
Total Nitrogen (µg/L)	910 - 910	910 (1)
Chlorophyll- uncorrected (µg/L)	1.7 - 1.7	1.7 (1)
Secchi (ft)	-	(0)
Secchi (m)	-	(0)
Color (Pt-Co Units)	188 - 188	188 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-8 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-8 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for CBA River-9 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	CBA River-9
Latitude	30.4676
Longitude	-86.0008
Water Body Type	Stream
Period of Record (year)	2008 to 2008
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for CBA River-9 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	7 - 7	7 (1)
Total Nitrogen (µg/L)	563 - 563	563 (1)
Chlorophyll- uncorrected (µg/L)	0.7 - 0.7	0.7 (1)
Secchi (ft)	3.0 - 3.0	3.0 (1)
Secchi (m)	0.9 - 0.9	0.9 (1)
Color (Pt-Co Units)	152 - 152	152 (1)
Specific Conductance (µS/cm@25 C)	-	(0)

LAKEWATCH Report for CBA River-9 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for CBA River-9 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Mullet Creek-1 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	Mullet Creek-1
Latitude	30.4797
Longitude	-86.3326
Water Body Type	Stream
Period of Record (year)	2002 to 2002
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for Mullet Creek-1 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	33 - 33	33 (1)
Total Nitrogen (µg/L)	380 - 380	380 (1)
Chlorophyll- uncorrected (µg/L)	5.0 - 5.0	5.0 (1)
Secchi (ft)	-	()
Secchi (m)	-	()
Color (Pt-Co Units)	-	()
Specific Conductance (µS/cm@25 C)	-	()

LAKEWATCH Report for Mullet Creek-1 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Mullet Creek-1 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Mullet Creek-2 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	Mullet Creek-2
Latitude	30.4803
Longitude	-86.3325
Water Body Type	Stream
Period of Record (year)	2002 to 2002
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for Mullet Creek-2 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	25 - 25	25 (1)
Total Nitrogen (µg/L)	280 - 280	280 (1)
Chlorophyll- uncorrected (µg/L)	1.0 - 1.0	1.0 (1)
Secchi (ft)	-	()
Secchi (m)	-	()
Color (Pt-Co Units)	-	()
Specific Conductance (µS/cm@25 C)	-	()

LAKEWATCH Report for Mullet Creek-2 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams

The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.

Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)				
Intercept (a)				
Slope (b)				
Coefficient of Determination (R ²)				
Probability of Significance (p)				
Potential Trend				

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Mullet Creek-2 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.

LAKEWATCH Report for Peach Creek-1 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	Peach Creek-1
Latitude	30.358
Longitude	-86.1079
Water Body Type	Stream
Period of Record (year)	1993 to 2016
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for Peach Creek-1 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	6 - 87	13 (24)
Total Nitrogen (µg/L)	300 - 680	465 (24)
Chlorophyll- uncorrected (µg/L)	0.7 - 5.6	2.1 (24)
Secchi (ft)	2.0 - 4.0	2.9 (24)
Secchi (m)	0.6 - 1.2	0.9 (24)
Color (Pt-Co Units)	66 - 271	161 (15)
Specific Conductance (µS/cm@25 C)	45 - 7554	2627 (10)

LAKEWATCH Report for Peach Creek-1 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	24	24	24	24
Intercept (a)	-841	-12507	25	45
Slope (b)	0.43	6.47	-0.01	-0.02
Coefficient of Determination (R ²)	0.03	0.23	0.00	0.09
Probability of Significance (p)	0.38	0.02	0.80	0.16
Potential Trend	No Trend	Increasing	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Peach Creek-1 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Peach Creek-1 (Walton)




Peach Creek-1 (Walton)



Peach Creek-1 (Walton)



Peach Creek-1 (Walton)



LAKEWATCH Report for Peach Creek-2 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	Peach Creek-2
Latitude	30.3576
Longitude	-86.1075
Water Body Type	Stream
Period of Record (year)	1993 to 2016
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for Peach Creek-2 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	5 - 107	13 (24)
Total Nitrogen (µg/L)	319 - 681	473 (24)
Chlorophyll- uncorrected (µg/L)	0.7 - 5.1	2.0 (24)
Secchi (ft)	2.2 - 6.3	3.3 (24)
Secchi (m)	0.7 - 1.9	1.0 (24)
Color (Pt-Co Units)	65 - 272	153 (15)
Specific Conductance (µS/cm@25 C)	62 - 11085	3178 (10)

LAKEWATCH Report for Peach Creek-2 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	24	24	24	24
Intercept (a)	-801	-12283	28	125
Slope (b)	0.41	6.36	-0.01	-0.06
Coefficient of Determination (R ²)	0.02	0.22	0.00	0.29
Probability of Significance (p)	0.51	0.02	0.76	0.01
Potential Trend	No Trend	Increasing	No Trend	Decreasing

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Peach Creek-2 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Peach Creek-2 (Walton)




Peach Creek-2 (Walton)



Peach Creek-2 (Walton)



Peach Creek-2 (Walton)



LAKEWATCH Report for Peach Creek-3 in Walton County Using Data Downloaded 10/17/2016

Introduction Streams

For many decades Florida has had a narrative nutrient water quality criterion in place to protect Florida's waters against nutrient over-enrichment. In 2009, the Florida Department of Environmental Protection (FDEP) initiated rulemaking and, by 2011, adopted what would be the first set of statewide numeric nutrient standards for Florida's waters. By 2015, almost all of the remaining waters in Florida have numeric nutrient standards (see for Florida Department of Environmental Regulation Nutrient Criteria's for: Streams, spring vents: <http://www.dep.state.fl.us/water/wqssp/nutrients/index.htm>).




Figure 1. Map showing nutrient thresholds areas for streams set forth by FDEP.

Base File Data: Definitions

The long-term data summary will include the following parameters listed with a definition after each one:

- **County:** Name of county in which the stream resides.
- **Name:** Stream name that LAKEWATCH uses for the system.
- **Latitude and Longitude:** Coordinates identifying the exact location of station 1 for each system.
- **Water Body Type:** Four different types of systems; lakes, estuaries, streams and springs.
- **Period of Record (year):** Years a stream has been in the LAKEWATCH program.

County	Walton
Name	Peach Creek-3
Latitude	30.3573
Longitude	-86.1072
Water Body Type	Stream
Period of Record (year)	1993 to 2016
Nutrient Watershed Region	Panhandle West

LAKEWATCH Report for Peach Creek-3 in Walton County Using Data Downloaded 10/17/2016

The nutrient thresholds for streams set forth by FDEP are listed in table below along with the map showing zones.

Nutrient Watershed Region	Total Phosphorus Nutrient Threshold ¹	Total Nitrogen Nutrient Threshold ¹
Panhandle West	60 µg/L	670 µg/L
Panhandle East	180 µg/L	1030 µg/L
North Central	300 µ/L	1870 µg/L
Peninsular	120 µg/L	1540 µg/L
West Central	490 µg/L	1650 µg/L
South Florida	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.	No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.

¹These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year periods.

Long-Term Data Summary Streams: Definitions

The following long-term data are the primary trophic state parameters collected by LAKEWATCH volunteers and classification variables color and specific conductance (LAKEWATCH recently began analyzing samples quarterly for color and specific conductance):

- **Total Phosphorus (µg/L):** The nutrient most often limiting growth of plant/algae in Florida's fresh and saltwater environments.
- **Total Nitrogen (µg/L):** Another nutrient needed for aquatic plant/algae growth but only limiting when nitrogen to phosphorus ratios are generally less than 10.
- **Chlorophyll-uncorrected (µg/L):** Chlorophyll concentrations are used to measure relative abundances of open water algal population.
- **Secchi (ft), Secchi (m):** Secchi measurements are estimates of water clarity (how far one can see into the water) and are listed with English and metric units.
- **Color (Pt-Co Units):** LAKEWATCH measures true color, which is the color of the water after particles have been filter out.
- **Specific Conductance (µS/cm@25°C):** Measurement of the ability of water to conduct electricity and can be used to estimate the amount of dissolve materials in water.

Long-Term Data Summary Streams: Data

Parameter	Minimum and Maximum Annual Means	Mean of Annual Means (Sampling years)
Total Phosphorus (µg/L)	5 - 86	12 (24)
Total Nitrogen (µg/L)	310 - 611	466 (24)
Chlorophyll- uncorrected (µg/L)	0.5 - 5.8	2.0 (24)
Secchi (ft)	2.1 - 3.9	3.0 (24)
Secchi (m)	0.6 - 1.2	0.9 (24)
Color (Pt-Co Units)	54 - 274	165 (15)
Specific Conductance (µS/cm@25 C)	44 - 7044	2536 (10)

LAKEWATCH Report for Peach Creek-3 in Walton County Using Data Downloaded 10/17/2016

Trend Analyses Streams


The following data are for linear regression statistics derived by plotting annual average total phosphorus, total nitrogen, chlorophyll, and Secchi data by year of data collection. Linear regression analysis is a common statistical approach used to determine if significant trends are occurring over time. These analyses define statistics based on the best fit line drawn through the data after plotting them with year on the horizontal line (x-axis) and the data value on the vertical line (y-axis). Figure 2 shows example plots with linear regression statistic of lakes that show significant total phosphorus increases, decreases and no change over time. The statistics that are listed include the following:

- **Number of years (n):** This is simply the number of years of data that were used to calculate annual means.
- **Intercept (a):** This is the value on the y-axis that the fitted line would have cross if the x-axis where zero.
- **Slope (b):** This is the rate at which the fitted line increases (positive number) or decreases (negative number).
- **Coefficient of determination (R²):** This value is an indication of how much variance above and below the fitted line there is in the data. This value ranges from 0 to 1. A high value means a tight fit and a low value means a loose fit.
- **Probability of Significance (p):** For most statistical analyses a p-value of less than 0.05 means the statistic is significant and analyses with p-values greater than 0.05 are not significant.


Statistic	Total Phosphorus	Total Nitrogen	Chlorophyll	Secchi
Number of Years (n)	24	24	24	24
Intercept (a)	-717	-13052	15	34
Slope (b)	0.36	6.74	-0.01	-0.02
Coefficient of Determination (R ²)	0.03	0.28	0.00	0.06
Probability of Significance (p)	0.45	0.01	0.90	0.26
Potential Trend	No Trend	Increasing	No Trend	No Trend

The following graphs on the next two pages are trend analyses examining regression between year and annual means of total phosphorus, total nitrogen, chlorophyll, and Secchi depth for Peach Creek-3 in Walton County. If there are no plots then there is less than five years of data, which is not enough for the analysis.


Peach Creek-3 (Walton)



Peach Creek-3 (Walton)



Peach Creek-3 (Walton)



Peach Creek-3 (Walton)

